arXiv:1304.1984v1 [cs.IT] 7 Apr 2013

A Multi-Dimensional Block-Circulant Perfect Array Construction

Samuel T. Blake
School of Mathematical Sciences, Monash University, Australia

Abstract. We present a N-dimensional generalization of the two-dimensional block-circulant perfect array construction by [Blake, 2013]. As in [Blake, 2013], the families of N-dimensional arrays possess pairwise *good* zero correlation zone (ZCZ) cross-correlation. Both constructions use a perfect autocorrelation sequence with the array orthogonality property (AOP).

This paper presents a generalization of the 2-dimensional block-circulant array construction by [Blake, 2013]. The generalized construction works in any number of dimensions, but is limited to the same size in each dimension as the original two-dimensional construction. Throughout the paper we follow the notation of [Blake, 2013].

We begin by stating the 2-dimensional construction for families of arrays with perfect autocorrelation and good cross-correlation as given in [Blake, 2013].

Construction I Let $\mathbf{a} = [a_0, a_1, \cdots, a_{n-1}]$ be a perfect sequence with the AOP for the divisor, d, and and $\mathbf{c} = [c(0), c(1), \cdots, c(d-1)]$ is a block of d perfect sequences – each of length m, where $m = 0 \bmod d$. We construct a family of arrays, $\mathbf{S}_k$, such that

$$\mathbf{S}_k = [S_{i,j}]_k = a_j \, c(j \bmod d)_{w\lfloor j/d \rfloor + k(j \bmod d) + i}$$

for $0 \le i < n$, $0 \le j < m$, $\mathbf{a}$ has the AOP for the divisor d, $0 < k \le m$, and $w = m/d$.

This construction produces perfect arrays up to size $r^2 \times r^2$ over r roots of unity. Each pair of distinct arrays has d^2 non-zero cross-correlation values, as $d \ll r$ we say the array has good ZCZ cross-correlation.

The generalized N-dimensional construction is given as follows.

Construction II Let $\mathbf{a} = [a_0, a_1, \cdots, a_{n-1}]$ be a perfect sequence with the AOP for the divisor, d, and $\mathbf{c} = [c(0), c(1), \cdots, c(d-1)]$ is a block of d perfect sequences – each of length m, where $m = 0 \bmod d$. We construct a family of k N-dimensional perfect arrays, $\mathbf{S}_k$, such that

$$\mathbf{S}_k = \left[S_{i_0, i_1, \cdots, i_{N-2}, j}\right]_k = a_j \prod_{v=0}^{N-2} c(j \bmod d)_{w\lfloor j/d \rfloor + k(j \bmod d) + i_v}$$

for $0 \le i_0, i_1, \cdots, i_{N-2} < n$, $0 \le j < m$, $0 < k \le m$, and $w = m/d$.

Each distinct pair of arrays from the family has d^2 non-zero cross-correlation values, regardless of the size of N. Thus, the ratio of zero to non-zero cross-correlation values is larger for higher dimensional arrays.

We show that $\mathbf{S}_k$ from Construction II has perfect autocorrelation ($k_1 = k_2$) and $\mathbf{S}_{k_1}$, $\mathbf{S}_{k_2}$ has *good* cross-correlation ($k_1 \neq k_2$). We now compute the cross-correlation of $\mathbf{S}_{k_1}$ with $\mathbf{S}_{k_2}$, $(0 < k_1, k_2 \le m)$ for shift $s_0, s_1, \cdots, s_{N-1}$:

$$
\begin{aligned}
&\theta_{\mathbf{S}_{k_1},\mathbf{S}_{k_2}}(s_0, s_1, \cdots, s_{N-1}) \\
&= \sum_{j=0}^{n-1}\sum_{i_0=0}^{m-1}\sum_{i_1=0}^{m-1}\cdots\sum_{i_{N-2}=0}^{m-1} \left[S_{i_0,i_1,\cdots,i_{N-2},j}\right]_{k_1}\left[S^*_{i_0+s_0,i_1+s_1,\cdots,j+s_{N-1}}\right]_{k_2} \\
&= \sum_{j=0}^{n-1}\sum_{i_0=0}^{m-1}\sum_{i_1=0}^{m-1}\cdots\sum_{i_{N-2}=0}^{m-1} \left(a_j \left(\prod_{v=0}^{N-2} c(j \bmod d)_{w\lfloor j/d\rfloor + k_1(j \bmod d)+i_v}\right) a^*_{j+s_{N-1}} \times \right. \\
&\qquad\qquad \left.\left(\prod_{v=0}^{N-2} c(j+s_{N-1} \bmod d)^*_{w\lfloor (j+s_{N-1})/d\rfloor + k_2(j+s_{N-1} \bmod d)+i_v+s_v}\right)\right) \\
&= \sum_{j=0}^{n-1}\left(a_j\, a^*_{j+s_{N-1}} \sum_{i_0=0}^{m-1}\sum_{i_1=0}^{m-1}\cdots\sum_{i_{N-2}=0}^{m-1}\left(\left(\prod_{v=0}^{N-2} c(j \bmod d)_{w\lfloor j/d\rfloor + k_1(j \bmod d)+i_v}\right)\times\right.\right. \\
&\qquad\qquad \left.\left.\left(\prod_{v=0}^{N-2} c(j+s_{N-1} \bmod d)^*_{w\lfloor (j+s_{N-1})/d\rfloor + k_2(j+s_{N-1} \bmod d)+i_v+s_v}\right)\right)\right) \qquad (7)
\end{aligned}
$$

Consider the case when $s_{N-1} = 0 \bmod d$. Let $s_{N-1} = l\,d$ and perform the change of coordinates $j = qd + r$, $(r < d)$. Then the inner summation in (7) becomes

$$
\begin{aligned}
&\sum_{i_0=0}^{m-1}\sum_{i_1=0}^{m-1}\cdots\sum_{i_{N-2}=0}^{m-1}\left(\left(\prod_{v=0}^{N-2} c(r)_{k_1 r+i_v}\right)\left(\prod_{v=0}^{N-2} c(r)^*_{lw+k_2 r+i_v+s_v}\right)\right) \\
&= \left(\sum_{i_0=0}^{m-1} c(r)_{k_1 r+i_0}\, c(r)^*_{lw+k_2 r+i_0+s_0}\right)\left(\sum_{i_1=0}^{m-1} c(r)_{k_1 r+i_1}\, c(r)^*_{lw+k_2 r+i_1+s_1}\right)\times \\
&\qquad \left(\sum_{i_2=0}^{m-1} c(r)_{k_1 r+i_2}\, c(r)^*_{lw+k_2 r+i_2+s_2}\right)\times\cdots\times\left(\sum_{i_{N-2}=0}^{m-1} c(r)_{k_1 r+i_{N-2}}\, c(r)^*_{lw+k_2 r+i_{N-2}+s_{N-2}}\right)
\end{aligned}
$$

which is independent of q. Thus, (7) becomes

$$
\begin{aligned}
&\sum_{r=0}^{d-1}\left(\left(\sum_{q=0}^{n/d-1} a_{qd+r}\, a^*_{qd+r+s_{N-1}}\right)\left(\sum_{i_0=0}^{m-1} c(r)_{k_1 r+i_0}\, c(r)^*_{lw+k_2 r+i_0+s_0}\right)\left(\sum_{i_1=0}^{m-1} c(r)_{k_1 r+i_1}\, c(r)^*_{lw+k_2 r+i_1+s_1}\right)\times\right. \\
&\qquad \left.\left(\sum_{i_2=0}^{m-1} c(r)_{k_1 r+i_2}\, c(r)^*_{lw+k_2 r+i_2+s_2}\right)\times\cdots\times\left(\sum_{i_{N-2}=0}^{m-1} c(r)_{k_1 r+i_{N-2}}\, c(r)^*_{lw+k_2 r+i_{N-2}+s_{N-2}}\right)\right) \qquad (8)
\end{aligned}
$$

As $\mathbf{a}$ has the AOP for the divisor d, the summation $\sum_{q=0}^{n/d-1} a_{qd+r}\, a^*_{qd+r+s_{N-1}} = 0$ for $s_{N-1} \neq 0 \bmod n/d$, thus $\theta_{\mathbf{S}_{k_1},\mathbf{S}_{k_2}}(s_0, s_1, \cdots, s_{N-1}) = 0$ for $s_{N-1} \neq 0 \bmod n/d$. Otherwise, for $s_{N-1} = 0 \bmod n/d$, (8) is zero when one of the remaining $N-2$ summations are zero. The summation over i_v is non-zero for $(k_2 - k_1)r + s_{N-1}m/d^2 + s_v = 0 \bmod m$. For each of the $s_{N-1} = 0, n/d, 2n/d, \cdots, (d-1)n/d$, there are d solutions for s_v, one for each $r = 0, 1, \cdots, d-1$. Thus, there are d^2 non-zero cross-correlation values for $s_{N-1} = 0 \bmod d$.

Now consider the case $k_1 = k_2$ (autocorrelation), (note that $s_{N-1} = 0 \bmod n/d$), then (8) becomes

$$\left(\sum_{r=0}^{d-1}\sum_{q=0}^{n/d-1} a_{qd+r}\, a^*_{qd+r+s_{N-1}}\right)\left(\sum_{i_0=0}^{m-1} c(r)_{i_0}\, c(r)^*_{lw+i_0+s_0}\right)\left(\sum_{i_1=0}^{m-1} c(r)_{i_1}\, c(r)^*_{lw+i_1+s_1}\right)\times \left(\sum_{i_2=0}^{m-1} c(r)_{i_2}\, c(r)^*_{lw+i_2+s_2}\right)\times\cdots\times\left(\sum_{i_{N-2}=0}^{m-1} c(r)_{i_{N-2}}\, c(r)^*_{lw+i_{N-2}+s_{N-2}}\right) \tag{8}$$

As $\mathbf{a}$ has the AOP for the divisor d, the left double summation is zero. Thus the autocorrelation is zero for $s_{N-1} = 0 \bmod d$.

Now, consider the case when $s_{N-1} \neq 0 \bmod d$. Let $s_{N-1} = l\,d + s$, $(s < d)$, and perform the change of coordinates $j = qd + r$, $(r < d)$. Then (7) becomes

$$\sum_{r=0}^{d-1}\sum_{q=0}^{n/d-1}\left(a_{qd+r}\, a^*_{qd+r+s_{N-1}} \sum_{i_0=0}^{m-1}\sum_{i_1=0}^{m-1}\cdots\sum_{i_{N-2}=0}^{m-1}\left(\left(\prod_{v=0}^{N-2} c(r)_{qw+k_1r+i_v}\right)\times \left(\prod_{v=0}^{N-2} c(r+s \bmod d)^*_{qw+lw+w\lfloor (r+s)/d\rfloor+k_2(r+s \bmod d)+i_v+s_v}\right)\right)\right)$$

$$= \sum_{r=0}^{d-1}\sum_{q=0}^{n/d-1}\left(a_{qd+r}\, a^*_{qd+r+s_{N-1}}\left(\sum_{i_0=0}^{m-1} c(r)_{qw+k_1r+i_0}\, c(r+s \bmod d)^*_{qw+lw+w\lfloor (r+s)/d\rfloor+k_2(r+s \bmod d)+i_0+s_0}\right)\times \left(\sum_{i_1=0}^{m-1} c(r)_{qw+k_1r+i_1}\, c(r+s \bmod d)^*_{qw+lw+w\lfloor (r+s)/d\rfloor+k_2(r+s \bmod d)+i_1+s_1}\right)\times\cdots\times \left(\sum_{i_{N-2}=0}^{m-1} c(r)_{qw+k_1r+i_{N-2}}\, c(r+s \bmod d)^*_{qw+lw+w\lfloor (r+s)/d\rfloor+k_2(r+s \bmod d)+i_{N-2}+s_{N-2}}\right)\right) \tag{9}$$

For each of the inner summations above, shifting both sequences qw places does not change the cross-correlation. So (9) becomes

$$= \sum_{r=0}^{d-1}\left(\left(\sum_{q=0}^{n/d-1} a_{qd+r}\, a^*_{qd+r+s_{N-1}}\right)\left(\sum_{i_0=0}^{m-1} c(r)_{k_1r+i_0}\, c(r+s \bmod d)^*_{lw+w\lfloor (r+s)/d\rfloor+k_2(r+s \bmod d)+i_0+s_0}\right)\times \left(\sum_{i_1=0}^{m-1} c(r)_{k_1r+i_1}\, c(r+s \bmod d)^*_{lw+w\lfloor (r+s)/d\rfloor+k_2(r+s \bmod d)+i_1+s_1}\right)\times\cdots\times \left(\sum_{i_{N-2}=0}^{m-1} c(r)_{k_1r+i_{N-2}}\, c(r+s \bmod d)^*_{lw+w\lfloor (r+s)/d\rfloor+k_2(r+s \bmod d)+i_{N-2}+s_{N-2}}\right)\right).$$

As the sequence $\mathbf{a}$ has the AOP, the summation

$$\sum_{q=0}^{n/d-1} a_{qd+r}\, a^*_{qd+r+s_{N-1}} = 0$$

for all r and s_{N-1} such that $s_{N-1} \neq 0 \bmod d$. Thus the cross-correlation of $\mathbf{S}_{k_1}$ with $\mathbf{S}_{k_2}$ is zero for $h \neq 0 \bmod d$ and all v. So Construction II produces a family of m perfect k-dimensional arrays with pairwise good cross-correlation.

It has been shown that perfect N-dimensional arrays can be constructed up to size $r^2 \times r^2 \times \cdots$ over r roots of unity. An exponential number of such arrays exist for each size. Furthermore, families of arrays possess good cross-correlation properties. Importantly, despite the size of N, the number of non-zero cross-correlation values is always d^2, so the ratio of the number of non-zero cross-correlation values to the number of entries in the arrays can be made as small as desired.

References

[Blake, 2013] S. Blake, T. E. Hall, A. Z. Tirkel, “Arrays over Roots of Unity with Perfect Autocorrelation and Good ZCZ Cross–Correlation”, accepted for publication, *Advances in Mathematics of Communications (AMC)*, 2013

Appendix I – Implementation of the Construction

In this appendix we give a Mathematica (version 8.0) implementation of the construction. (Arrays are given in index notation, that is, the mapping: $e^{2\pi\sqrt{-1}s_n/r} \rightarrow s_n$.)

We begin with the code for periodic cross-correlation, `XCV` and autocorrelation, `ACV`:

```
In[1]:= XCV[a_, b_, r_Integer] := Block[{A, B},
  A = Developer`ToPackedArray[Exp[(2. Pi I a)/r]];
  B = Developer`ToPackedArray[Exp[(-2. Pi I b)/r]];
  Chop[ListCorrelate[A, B, 1], 1*^-5]]

In[2]:= ACV[m_, r_Integer] := XCV[m, m, r]
```

The N-dimensional array construction, `ConstructionND`, takes as input the multiplying sequence, **a**, the block of d perfect sequences, **c**, the parameter, k, the number of dimensions, *dims*, and the number of roots of unity

```
In[3]:= aref = #1[[#2 + 1]] &;

In[4]:= ConstructionND[a_, c_, k_, dims_] :=
 With[{d = Length[c], l = Length[First[c]]},
    Table[
       Array[
          indexND[a, c, k, d, l, j, ##] &, Table[l, {dims - 1}], 0],
   {j, 0, Length[a] - 1}]]

In[5]:= indexND[a_, c_, k_, d_, l_, index__] :=
 With[{w = Length[First[c]]/d, j = First @ {index}},
     aref[a, j] + Product[aref[aref[c, Mod[j, d]],
        Mod[w Floor[j/d] + k Mod[j, d] + i, l]], {i, Rest @ {index}}]]
```

We construct a perfect 4 × 4 × 4 × 4 binary array and compute its periodic autocorrelation:

```
In[6]:= ConstructionND[FrankSequence[2], {#, Decimate[#, 3]} & @ FrankSequence[2], 0, 4];
Mod[%, 2]
Count[ACV[%, 2], n_?NumericQ /; n != 0., Infinity]
```

```
        ⎡ 0 0 0 1   0 0 0 1   0 0 0 1   1 1 1 0 ⎤
        ⎢ 0 0 0 1   0 0 0 1   0 0 0 1   1 1 1 0 ⎥
        ⎢ 0 0 0 1   0 0 0 1   0 0 0 1   1 1 1 0 ⎥
        ⎢ 1 1 1 0   1 1 1 0   1 1 1 0   0 0 0 1 ⎥
        ⎢ 0 1 0 0   1 0 1 1   0 1 0 0   0 1 0 0 ⎥
        ⎢ 1 0 1 1   0 1 0 0   1 0 1 1   1 0 1 1 ⎥
        ⎢ 0 1 0 0   1 0 1 1   0 1 0 0   0 1 0 0 ⎥
Out[7]= ⎢ 0 1 0 0   1 0 1 1   0 1 0 0   0 1 0 0 ⎥
        ⎢ 0 1 0 0   1 0 1 1   0 1 0 0   0 1 0 0 ⎥
        ⎢ 1 0 1 1   0 1 0 0   1 0 1 1   1 0 1 1 ⎥
        ⎢ 0 1 0 0   1 0 1 1   0 1 0 0   0 1 0 0 ⎥
        ⎢ 0 1 0 0   1 0 1 1   0 1 0 0   0 1 0 0 ⎥
        ⎢ 1 1 1 0   1 1 1 0   1 1 1 0   0 0 0 1 ⎥
        ⎢ 1 1 1 0   1 1 1 0   1 1 1 0   0 0 0 1 ⎥
        ⎢ 1 1 1 0   1 1 1 0   1 1 1 0   0 0 0 1 ⎥
        ⎣ 0 0 0 1   0 0 0 1   0 0 0 1   1 1 1 0 ⎦
```

```
Out[8]= 1
```

The following generalization of `indexND` allows for computations over the quaternions:

```
In[9]:= NonCommutativeDot[e_] := e
```

```
In[10]:= indexND[a_, c_, k_, d_, l_, index__] /; ! FreeQ[a, Quaternion] :=
 With[{w = Length[First[c]]/d, j = First @ {index}},
  aref[a, j] ** Apply[NonCommutativeDot,
    Table[aref[aref[c, Mod[j, d]], Mod[w Floor[j/d] + k Mod[j, d] + i, l]], {i, Rest@{index}}]]]
```

The following is a perfect quaternion sequence which posesses the array orthogonality property for the divisor 4:

$$[1, k, 1, -k, -i, -k, i, -k, -1, k, -1, -k, i, -k, -i, -k]$$

We use this sequence to construct a perfect 16×16 quaternion array.

```
In[11]:= q16 = {Quaternion[1, 0, 0, 0], Quaternion[0, 0, 0, 1],
   Quaternion[1, 0, 0, 0], Quaternion[0, 0, 0, -1],
   Quaternion[0, -1, 0, 0], Quaternion[0, 0, 0, -1],
   Quaternion[0, 1, 0, 0], Quaternion[0, 0, 0, -1],
   Quaternion[-1, 0, 0, 0], Quaternion[0, 0, 0, 1],
   Quaternion[-1, 0, 0, 0], Quaternion[0, 0, 0, -1],
   Quaternion[0, 1, 0, 0], Quaternion[0, 0, 0, -1],
   Quaternion[0, -1, 0, 0], Quaternion[0, 0, 0, -1]};
```

```
In[12]:= ConstructionND[q16, {#, Decimate[#, 3], RotateRight[#, 2], #} & @ q16, 0, 2]
Count[QuaternionCV2D[%], n_?NumericQ /; n != 0., Infinity]
```

`Out[12]=`

$$\begin{bmatrix}
1 & k & 1 & -k & -i & -k & i & -k & -1 & k & -1 & -k & i & -k & -i & -k \\
k & 1 & j & -1 & j & 1 & k & 1 & -k & 1 & -j & -1 & -j & 1 & -k & 1 \\
-i & -k & 1 & k & 1 & -k & -i & -k & i & -k & -1 & k & -1 & -k & i & -k \\
-k & 1 & -k & -1 & j & -1 & -j & -1 & k & 1 & k & -1 & -j & -1 & j & -1 \\
-1 & -j & 1 & -j & i & j & i & -j & 1 & -j & -1 & -j & -i & j & -i & -j \\
-j & -1 & -k & -1 & k & -1 & j & 1 & j & -1 & k & -1 & -k & -1 & -j & 1 \\
i & j & 1 & j & -1 & j & -i & -j & -i & j & -1 & j & 1 & j & i & -j \\
j & -1 & -j & -1 & k & 1 & k & -1 & -j & -1 & j & -1 & -k & 1 & -k & -1 \\
1 & -k & 1 & k & -i & k & i & k & -1 & -k & -1 & k & i & k & -i & k \\
-k & 1 & -j & -1 & -j & 1 & -k & 1 & k & 1 & j & -1 & j & 1 & k & 1 \\
-i & k & 1 & -k & 1 & k & -i & k & i & k & -1 & -k & -1 & k & i & k \\
k & 1 & k & -1 & -j & -1 & j & -1 & -k & 1 & -k & -1 & j & -1 & -j & -1 \\
-1 & j & 1 & j & i & -j & i & j & 1 & j & -1 & j & -i & -j & -i & j \\
j & -1 & k & -1 & -k & -1 & -j & 1 & -j & -1 & -k & -1 & k & -1 & j & 1 \\
i & -j & 1 & -j & -1 & -j & -i & j & -i & -j & -1 & -j & 1 & -j & i & j \\
-j & -1 & j & -1 & -k & 1 & -k & -1 & j & -1 & -j & -1 & k & 1 & k & -1
\end{bmatrix}$$

`Out[13]=` 1

We now construct a family of 9 perfect $9 \times 9 \times 9 \times 9$ arrays over 3 roots of unity. The perfect Frank sequence: $[0, 0, 0, 0, 1, 2, 0, 2, 1]$, is used as the base sequence.

```
In[14]:= family3D = Table[
   ConstructionND[FrankSequence[3],
      {Decimate[#, 2], Decimate[#, 5], Decimate[#, 7]} & @ FrankSequence[3], n, 4],
  {n, 9}];
```

Each pair of sequences from the family have $d^2 = 3^2$ non-zero cross-correlation values. Here's the non-zero cross-correlation values of an example pair:

```
In[15]:= XCV[family3D[[1]], family3D[[2]], 3];
Cases[%, n_?NumericQ /; n != 0., Infinity]

Out[16]= {2187., 2187., 2187., -1093.5 - 1894. I, -1093.5 + 1894. I, 2187., -1093.5 + 1894. I,
-1093.5 - 1894. I, 2187.}
```

Here we generate all pairs of arrays from the family and count the number of non-zero cross-correlation values for each pair.

```
In[17]:= Count[XCV[#1, #2, 3], n_?NumericQ /; n != 0., Infinity] & @@@ Tuples[family3D, 2]

Out[17]= {1, 9, 9, 9, 9, 9, 9, 9, 9, 9, 1, 9, 9, 9, 9, 9, 9, 9, 9, 9, 1, 9, 9, 9, 9, 9, 9, 9, 9,
9, 1, 9, 9, 9, 9, 9, 9, 9, 9, 9, 1, 9, 9, 9, 9, 9, 9, 9, 9, 9, 1, 9, 9, 9, 9, 9, 9, 9, 9, 9, 1,
9, 9, 9, 9, 9, 9, 9, 9, 9, 1, 9, 9, 9, 9, 9, 9, 9, 9, 9, 1}
```

(The 1's above correspond to the autocorrelation of non distinct pairs.)